

RESULTS

We demonstrated that AGF is expressed in proopiomelanocortin (POMC)-positive neurons located in the arcuate nucleus (ARC) of the hypothalamus. AGF expression was stimulated by leptin-induced STAT3 phosphorylation. Notably, intracerebroventricular injection of AGF significantly reduced food intake by stimulating phosphorylation of CREB in the POMC and increasing α -melanocyte-stimulating hormone (α -MSH) content in the hypothalamus. We also found that hypothalamic injection of AGF significantly suppressed food intake and decreased body weight in high-fat-diet-induced obese mice, which exhibit leptin insensitivity.

CONCLUSION

Collectively, our findings demonstrate that hypothalamic AGF provokes the anorectic melanocortin pathway and mediates leptin signaling to prevent obesity.

KEY WORDS

Obesity, AGF, Hypothalamus, Leptin signaling

OA-GE-09

DETERMINANTS OF INSULIN RESISTANCE AMONG PEOPLE LIVING IN A RURAL AREA OF INDONESIA

<https://doi.org/10.15605/jafes.034.02.S71>

Dicky Levenus Tahapary,^{1,2} Dante Saksono Harbuwono,^{1,2} Pradana Soewondo,^{1,2}

¹*Division of Metabolism and Endocrinology, Department of Internal Medicine, Dr. Cipto Mangunkusumo National General Hospital, Faculty of Medicine Universitas Indonesia*

²*Metabolic, Cardiovascular and Aging Cluster, The Indonesian Medical Education and Research Institute, Faculty of Medicine Universitas Indonesia*

INTRODUCTION

Worldwide prevalence of obesity has increased rapidly, of which the main driver is the rising of body weight among people living in the rural area. While obesity is one of the main risk factors for the development of type 2 diabetes (T2DM), in a rural area, other environmental factors might also play a role. This study aims to assess the determinants of insulin resistance (IR), the main pathogenesis of T2DM, among people living in a rural area of Indonesia.

METHODOLOGY

We analyzed the baseline data of the Sugarspin study (n=1669) which was conducted in Nangapanda, Flores Island, Indonesia. The influence of each risk factors to HOMA-IR, was analyzed using linear regression analysis.

RESULTS

The prevalence of IR was 27.8% and 28.2% for men and women respectively. Increasing BMI and waist circumference were associated with increasing IR. Increasing age was also associated with increasing IR, however, in men after the age of 55, further increment in age was not associated with increasing IR. Progressive increase in BMI was observed until the age of 40, which then plateau until the age of 55, and then decline afterward. Interestingly, helminth infection was independently associated with a lower IR.

CONCLUSION

In a rural area, the main determinant of IR was similar to the one we observed in an urban area, which is obesity. However, the more prevalent infectious disease in a rural area might also directly or indirectly influence IR by its impact on obesity or other factors which need to be elucidated further.

KEY WORDS

insulin resistance, determinants, Indonesia, obesity, environments, rural

OA-GE-10

ASSOCIATION OF BODY MASS INDEX AND ALL CAUSE MORTALITY IN A TERTIARY REGIONAL HOSPITAL

<https://doi.org/10.15605/jafes.034.02.S72>

Lei Anne Carolino, Maria Luisa Arkoncel, Misael Cruz

Jose B. Lingad Memorial Regional Hospital, San Fernando, Pampanga, Philippines

INTRODUCTION

This study is conducted to provide further clinical evidence to determine whether there is a significant association between body mass index and all-cause mortality risk. At present, there are no existing studies done in the Philippines regarding exploring BMI as risk factor for mortality among patients with existing illnesses.

METHODOLOGY

This prospective cohort study was conducted from September 2016 until September 2017. Analysis of BMI and mortality was done and logistic regression was performed to determine confounding variables.