

## Before Low Serum Cortisol Can Be Attributed to SARS-CoV-2 Infection, Alternative Causes Must Be Ruled Out

Josef Finsterer,<sup>1</sup> ArunSundar MohanaSundaram,<sup>2</sup> Sounira Mehri<sup>3</sup>

<sup>1</sup>Neurology and Neurophysiology Center, Vienna, Austria

<sup>2</sup>School of Pharmacy, Sathyabama Institute of Science and Technology, Chennai, India,

<sup>3</sup>Biochemistry Laboratory, Faculty of Medicine, Monastir, Tunisia

**Key words:** SARS-CoV-2 infection, cortisol, ACTH, adrenal insufficiency, COVID-19

We read with interest Banu et al.'s article about a cross-sectional study of adrenal function using blood levels of cortisol and adrenocorticotropic hormone (ACTH) in 91 patients with acute SARS-CoV-2 infection. It was found that 27% of patients with severe COVID-19, 26% with mild COVID-19 and 4% with moderate COVID-19 had adrenal insufficiency. Decreased cortisol reserve was found in 6.6% of all patients. It was concluded that the adrenocortical response was impaired in a significant number of non-critically ill COVID-19 patients, with the percentage being highest in patients with severe COVID-19 disease.<sup>1</sup> The study is impressive, but some points should be discussed.

The major limitation of the study is that imaging of the pituitary gland and hypothalamus was not performed. To rule out pituitary adenoma, hypophysitis, pituitary apoplexy, pituitary abscess, sellar tuberculoma, or hypothalamic stroke or bleeding, magnetic resonance imaging (MRI) of the pituitary gland and the hypothalamus with contrast agent must be performed. Like any other infection, SARS-CoV-2 infection can trigger the appearance of symptoms of a pituitary pathology.<sup>2</sup>

A second limitation is that imaging of the suprarenal glands was not performed. To assess whether there was adrenal atrophy, adenoma, ischemia, bleeding, or carcinoma, adrenal imaging is mandatory.<sup>3</sup>

A third limitation is that current medication was not reported. Several medications are known to affect serum

cortisol levels, making it imperative to know all medications regularly taken by the included patients. Ketoconazole, isilodrostat, vitamin D and omega-3 fatty acids in particular are known to be able to reduce cortisol levels in the blood.<sup>4</sup> A fourth limitation is that no ACTH stimulation test was performed to determine whether primary or secondary suprarenal insufficiency was present. Determining cortisol levels alone without ACTH stimulation testing is not reliable with regard to cortical adrenal function.

A fifth limitation is that serum dehydroepiandrosterone (DHEA) levels were not measured. Since DHEA is the antagonist of cortisol and decreases with age, it is conceivable that high levels of DHEA promoted the reduction of serum cortisol.<sup>5</sup>

Since low thyroxine levels may be associated with low cortisol levels,<sup>6</sup> it would also be desirable to know whether thyroid function was normal or abnormal in the included patients.

In summary, the excellent study has limitations that should be addressed before drawing conclusions. Clarifying the weaknesses would strengthen the conclusions and could improve the study. Serum cortisol levels may be low not only due to SARS-CoV-2 infection but also due to several other causes. Before adrenal insufficiency can be attributed solely to inadequate cortical cortisol production, the entire hypothalamic-pituitary-adrenal axis must be thoroughly examined.

**Statement of Authorship**

All authors certified fulfillment of ICMJE authorship criteria.

**CRedit Author Statement**

**JF:** Conceptualization, Software, Validation, Formal analysis, Investigation, Resources, Data curation, Writing – original draft preparation, Writing – review and editing, Supervision, Project administration, Funding acquisition; **ASMS:** Methodology, Investigation, Writing – review and editing, Supervision; **SM:** Methodology, Investigation, Writing – review and editing

**Author Disclosure**

The authors declared no conflict of interest.

**Data Availability Statement**

No datasets were generated or analyzed for this study.

**Funding Source**

None.

**References**

1. Banu H, Sultana N, Md Shahed M, Hasanat MA, Saleh AA, Arafat SM. Hypothalamic-pituitary-adrenal axis activity in SARS-CoV-2 infected noncritically ill hospitalized patients. *J ASEAN Fed Endocr Soc.* 2023;38(2):65-70. PMID: 38045674 PMID: 38045674 PMID: 38045674 DOI: 10.15605/jafes.038.02.04
2. Rao S, Rajkumar A, Kuruvilla S. Sellar lesion: not always a pituitary adenoma. *Indian J Pathol Microbiol.* 2008;51(2):269-70. PMID: 18603706 DOI: 10.4103/0377-4929.41688
3. Leyendecker P, Ritter S, Riou M, et al. Acute adrenal infarction as an incidental CT finding and a potential prognosis factor in severe SARS-CoV-2 infection: A retrospective cohort analysis on 219 patients. *Eur Radiol.* 2021;31(2):895-900. PMID: 32852586 PMID: 32852586 PMID: 32852586 DOI: 10.1007/s00330-020-07226-5
4. Amodru V, Brue T, Castinetti F. Synergistic cortisol suppression by ketoconazole-osilodrostat combination therapy. *Endocrinol Diabetes Metab Case Rep.* 2021;2021:21-0071. PMID: 34877930 PMID: 34877930 PMID: 34877930 DOI: 10.1530/EDM-21-007
5. Hechter O, Grossman A, Chatterton RT Jr. Relationship of dehydroepiandrosterone and cortisol in disease. *Med Hypotheses.* 1997;49(1):85-91. PMID: 9247914 DOI: 10.1016/s0306-9877(97)90258-9
6. Mazur A. Do cortisol and thyroxin correlate with nervousness and depression among male army veterans? *Biol Psychol.* 1994;37(3):259-63. PMID: 7948470 DOI: 10.1016/0301-0511(94)90007-8.

Authors are required to accomplish, sign and submit scanned copies of the JAFES Author Form consisting of: (1) Authorship Certification, that authors contributed substantially to the work, that the manuscript has been read and approved by all authors, and that the requirements for authorship have been met by each author; (2) the Author Declaration, that the article represents original material that is not being considered for publication or has not been published or accepted for publication elsewhere, that the article does not infringe or violate any copyrights or intellectual property rights, and that no references have been made to predatory/suspected predatory journals; (3) the Author Contribution Disclosure, which lists the specific contributions of authors; (4) the Author Publishing Agreement which retains author copyright, grants publishing and distribution rights to JAFES, and allows JAFES to apply and enforce an Attribution-Non-Commercial Creative Commons user license; and (5) the Conversion to Visual Abstracts (\*optional for original articles only) to improve dissemination to practitioners and lay readers. Authors are also required to accomplish, sign, and submit the signed ICMJE form for Disclosure of Potential Conflicts of Interest. For original articles, authors are required to submit a scanned copy of the Ethics Review Approval of their research as well as registration in trial registries as appropriate. For manuscripts reporting data from studies involving animals, authors are required to submit a scanned copy of the Institutional Animal Care and Use Committee approval. For Case Reports or Series, and Images in Endocrinology, consent forms, are required for the publication of information about patients; otherwise, appropriate ethical clearance has been obtained from the institutional review board. Articles and any other material published in the JAFES represent the work of the author(s) and should not be construed to reflect the opinions of the Editors or the Publisher.



Experience the new JAFES.  
Visit us at [www.ASEAN-endocrinejournal.org](http://www.ASEAN-endocrinejournal.org).