

## Ethnic Disparity in Inter-Arm Systolic Blood Pressure Difference and its Determinants among Asians with Type 2 Diabetes: A Cross-Sectional Study

Xiao Zhang,<sup>1</sup> Jian Jun Liu,<sup>1</sup> Chee Fang Sum,<sup>2</sup> Yeoh Lee Ying,<sup>3</sup> Subramaniam Tavintharan,<sup>2</sup> Na Li,<sup>1</sup> Chang Su,<sup>1</sup> Serena Low,<sup>1</sup> Simon BM Lee,<sup>4</sup> Wern Ee Tang,<sup>4</sup> Su Chi Lim<sup>2</sup>

<sup>1</sup>Clinical Research Unit, Khoo Teck Puat Hospital, Singapore

<sup>2</sup>Diabetes Centre and Department of Medicine, Khoo Teck Puat Hospital, Singapore

<sup>3</sup>Department of Medicine, Khoo Teck Puat Hospital, Singapore

<sup>4</sup>National Healthcare Group Polyclinics, Singapore

### Abstract

**Objectives.** An inter-arm difference in systolic blood pressure (IADSBP) of 10 mmHg or more has been associated with cardiovascular disease (CVD) and increased mortality in T2DM patients. We aim to study ethnic disparity in IADSBP and its determinants in a multi-ethnic T2DM Asian cohort.

**Methodology.** Bilateral blood pressures were collected sequentially in Chinese (n=654), Malays (n=266) and Indians (n=313). IADSBP was analyzed as categories (<10 and ≥10 mmHg). Logistic regression model was used to evaluate the determinants of IADSBP ≥10 mmHg.

**Results.** Malays (27.4%) and Indians (22.4%) had higher prevalence of IADSBP ≥10 mmHg than Chinese (17.4%) (p=0.002). After adjustment for age, gender, duration of diabetes, hemoglobin A1c, body mass index (BMI), heart rate, pulse wave velocity (PWV), estimated glomerular filtration rate (eGFR), albumin-to-creatinine ratio (ACR), smoking, hypertension, soluble receptor for advanced glycation end products (sRAGE), and usage of hypertension medications, ethnicity remained associated with IADSBP. While Malays were more likely to have IADSBP ≥10 mmHg than Chinese (OR=1.648, 95%CI: 1.138-2.400, p=0.009), Indians had comparable odds with the Chinese. BMI (OR=1.054, 95%CI: 1.022-1.087, p=0.001) and hypertension (OR=2.529, 95%CI: 1.811-3.533, p<0.001) were also associated with IADSBP ≥10 mmHg.

**Conclusion.** IADSBP in Malays were more likely to be ≥10 mmHg than the Chinese which may explain their higher risk for CVD and mortality. Measuring bilateral blood pressures may identify high-risk T2DM individuals for intensive risk factor-management.

**Key words:** type 2 diabetes, cardiovascular disease, inter-arm difference in systolic blood pressure

### INTRODUCTION

Type 2 diabetes (T2DM) is a rapidly evolving global health issue<sup>1</sup> and Asia is the epicenter of this worldwide epidemic.<sup>2</sup> The prevalence of T2DM has been predicted to double from 7.3% in 1990 to 15% in 2050 in Singapore, a multi-ethnic city-state composed of three major ethnic groups (Chinese, Malays and Indians).<sup>3</sup> The ethnic diversity in Singapore provided us a unique opportunity to study the ethnic disparity on T2DM-related adverse outcomes.

Ethnic disparity in the development of diabetic complications, such as cardiovascular disease (CVD) and mortality from CVD, has been reported in many Western countries.<sup>4,5</sup> Such ethnic disparity has also been found

among Asians with T2DM in Singapore, showing higher risk for ischemic heart disease and cardiovascular mortality in Indians and Malays compared to Chinese.<sup>6-9</sup> In our recent studies in Singapore, we also revealed higher risk for diabetic complications in Indians and Malays compared to Chinese, such as acute myocardial infarction and cardiovascular mortality.<sup>10,11</sup> Several reasons have been suggested to explain their excess adverse outcomes, including less favorable metabolic and lipid profiles e.g., body mass index (BMI); lifestyle, e.g., less physical activity, less fruit and vegetable consumption; and poorer diabetes control, e.g., higher hemoglobin A1c (HbA1c). Furthermore, we observed higher central arterial stiffness, a well-established biomarker for vasculopathy, and lower level of soluble receptor for advanced glycation end-

eISSN 2308-118x

Printed in the Philippines

Copyright © 2016 by the JAFES

Received: April 22, 2016. Accepted: July 13, 2016.

<https://doi.org/10.15605/jafes.031.02.02>

Corresponding author: Assoc. Prof. Su Chi Lim, MBBS, MRCP, PhD  
Senior Consultant

Khoo Teck Puat Hospital

90 Yishun Central, Singapore 768828

Tel. No.: 65-66922353

Fax No.: 65-66023772

E-mail: [lim.su.chi@alexandrahealth.com.sg](mailto:lim.su.chi@alexandrahealth.com.sg)

products (sRAGE), a molecule with protective function on the vasculature in Malays and Indians than Chinese.<sup>12</sup> However, the mechanisms underlying the ethnic disparity in multifactorial diseases, such as CVD still remain largely unknown.

An inter-arm difference in blood pressure of less than 10 mmHg is classified as normal based on the new clinical guideline for hypertension.<sup>13</sup> Evolving evidence suggested that an inter-arm difference in systolic blood pressure (IADSBP) of 10 mmHg or more is associated with CVD and cardiovascular mortality.<sup>14,15</sup> It is interesting that the prevalence of IADSBP  $\geq$ 10 mmHg is higher among T2DM patients (~9-10%) when compared to the general population (4.4%).<sup>16,17</sup> Recently, IADSBP  $\geq$ 10 mmHg has been associated with CVD and cardiovascular mortality in T2DM patients, suggesting that the detection of IADSBP may identify a subpopulation at high risk of vascular events among T2DM patients.<sup>18</sup>

The prevalence of IADSBP  $\geq$ 10 mmHg varies widely across different studies, depending on ethnicity and background of enrolled individuals.<sup>14,15</sup> For example, the prevalence of IADSBP  $\geq$ 10 mmHg or the mean of IADSBP in Asians<sup>19-21</sup> is generally lower than that in Caucasians.<sup>18,22,23</sup> In a multi-ethnic study of atherosclerosis in USA, different prevalence rates of IADSBP  $\geq$ 15 mmHg were observed in African-Americans, Caucasians and Chinese.<sup>24</sup> However, disaggregated information on IADSBP is limited among Asian sub-ethnicities with T2DM, notwithstanding the existence of clear ethnic disparity in diabetes-related adverse outcomes. Therefore, we aim to investigate ethnic differences in IADSBP in a multi-ethnic T2DM Asian cohort.

## METHODOLOGY

### Study population and design

The Singapore Study of Macro-angiopathy and Micro-vascular Reactivity in Type 2 Diabetes (SMART2D) is a cross-sectional study conducted between August 2011 and February 2014 which included a total of 2,057 adults aged 21–90 years with T2DM. Inclusion and exclusion criteria of SMART2D have been previously described.<sup>25</sup> We included 1,233 individuals with bilateral BP measurements in this study. These selected individuals have similar profile compared to the rest of the subjects (detailed data not shown). This study has been approved by our institution's domain-specific ethics review committee. Individual written informed consent was obtained prior to enrollment in the study.

### Clinical and biochemical measurements

BMI was calculated as body weight (kg)/height (m)<sup>2</sup>. Urinary albumin-to-creatinine ratio (ACR) was determined by urinary creatinine measured by enzymatic

method on Roche/Hitachi cobas c system (Roche Diagnostic GmbH, Mannheim, Germany) and albumin measured by a solid-phase competitive chemiluminescent enzymatic immunoassay with a lower detection limit of 2.5  $\mu$ g/ml (Immulite; DPC, Gwynedd, UK). Estimated glomerular filtration rate (eGFR) was calculated based on a widely used Modified Diet in Renal Disease (MDRD) equation in patients with diabetes.<sup>26</sup> HbA1c was measured based on monoclonal antibody agglutination reaction using a point-of-care immunoassay analyzer (DCA Vantage Analyzer; Siemens, Erlangen, Germany) certified by National Glycohemoglobin Standardization Program. High-density lipoprotein-cholesterol (HDL-C) and low-density lipoprotein-cholesterol (LDL-C) were quantified by enzymatic method using Kodak Ektachem chemistry slides. Total triglyceride was quantified by enzymatic colorimetric method on Roches/Hitachi cobas c system. Carotid-femoral pulse wave velocity (PWV) was measured by the foot-to-foot method using a well-validated device, SphygmoCor® (AtCor Medical, Sydney, Australia) as described previously.<sup>27</sup> PWV was expressed as the distance between the two recording sites (meters) divided by transit time (seconds). Total sRAGE was quantified by ELISA (R&D Systems, Minneapolis, MN) according to manufacturer's protocol. The intra- and inter-assay coefficients of variation were 5.7% and 7.7%, respectively. The sensitivity reported by the manufacturer was 4.12 pg/ml.

### Blood pressure measurement

Sequentially repeated measurements of bilateral blood pressures were performed using mercury sphygmomanometers, the "gold standard" device for blood pressure measurement.<sup>28</sup> An appropriately sized cuff is attached to the upper arm, and a series of calibration readings are taken with the mercury sphygmomanometer to ensure that the device is giving accurate readings. After five minutes of seated rest, sphygmomanometer was placed on the arm, alternating right then left, and allowing one minute between each measurement on the same arm. Systolic blood pressure (SBP) and diastolic blood pressure (DBP) were calculated from the average of three most consistent readings. At the end of the five-minute rest, heart rate was assessed by an OMRON® digital blood pressure monitor.

### Statistical analysis

Standard descriptive statistics were used to summarize the characteristics of individuals with T2DM. Normally distributed continuous data were expressed as means and standard deviations (SDs), and skewed variables were expressed as median and inter-quartile range and nature log (ln)-transformed before data analysis. Differences among ethnicity were compared by one-way ANOVA or Kruskal–Wallis test where appropriate.

IADSBP was analyzed as categories (<10 and ≥10 mmHg). Univariate logistic regression model was used to evaluate the determinants of IADSBP ≥10 mmHg. Variables that are statistically significant in univariate analysis (p<0.25)<sup>29</sup> (Table S1) or known to potentially affect IADSBP and PWV<sup>15,30-32</sup> were added into the model. Age, gender, ethnicity, duration of diabetes, HbA1c, BMI, heart rate, PWV, smoking, hypertension (defined as any SBP ≥140 mmHg or DBP ≥90 mmHg<sup>33</sup>), eGFR, ACR, sRAGE, and commonly used medications in diabetes e.g., angiotensin-converting-enzyme (ACE) inhibitors and angiotensin receptor blockers (ARBs), were added into multivariable logistic regression model. All statistical analyses were performed using IBM SPSS (Version 22). A two-tailed p value of less than 0.05 was considered as statistically significant.

In collinearity test for each variable, the tolerance value was >0.2 and variance inflation factor was <3 for all variables, indicating that there was no serious problem with multicollinearity.

**RESULTS**

Table 1 shows the characteristics of T2DM patients stratified by ethnicity, including 654 Chinese (53.0%), 266 Malays (21.6%) and 313 Indians (25.4%). Compared to Chinese, Malays and Indians were younger and had higher BMI. Malays had higher heart rate, HbA1c and ACR, and lower eGFR than Chinese and Indians. The percentages of ischemic heart disease, coronary balloon angioplasty, and history of myocardial infarction in Indians are higher than

Chinese and Malays. There were no significant differences in percentages of current smoker, coronary bypass operation, stroke and proliferative retinopathy in Malays compared with Indians and Chinese.

Compared to Chinese (5.4±4.8 mmHg), IADSBP is significantly higher in Malays (7.1±6.4 mmHg, p<0.001) and non-significantly higher in Indians (6.1±6.0 mmHg, p=0.132). There is significant higher percentage of IADSBP ≥10 mmHg in Malays (27.4%) and Indians (22.4%) than Chinese (17.4%, P=0.002).

Univariate analysis shows a higher odds for IADSBP ≥10 mmHg in Malays (OR=1.791, 95% CI: 1.279-2.508, p=0.001) than the Chinese and comparable odds of Indians with the Chinese (OR=1.364, 95% CI: 0.997-1.905, p=0.068). Table 2 shows the associations of IADSBP ≥10 mmHg with ethnicity, and clinical and biochemical determinants by multivariable logistic regression models. After adjustment for age, gender, duration of diabetes, hemoglobin A1c, BMI, heart rate, PWV, eGFR, ACR, smoking, hypertension, sRAGE, and usage of hypertension medications, ethnicity remains significantly associated with IADSBP. While odds for Indians are comparable to Chinese (p=0.089), Malays are more likely to have elevated IADSBP of at least 10 mmHg (OR=1.648, 95% CI: 1.138-2.400, p=0.009). BMI (OR=1.054, 95%CI: 1.002-1.087, p=0.001) and hypertension (OR=2.529, 95%CI: 1.811-3.533), p<0.001) are also significant determinants of IADSBP ≥10 mmHg.

**Table 1.** Clinical and biochemical characteristics of individuals with T2DM stratified by ethnicity (n=1,233)

Variables	Total (1,233)	Malay (266)	Indian (313)	Chinese (654)	P-value
Entry age (yrs)	57.7±10.4	55.9±9.9	56.2±9.7	59.2±10.8	<0.001
Male gender (%)	51.8	45.8	50.2	55.0	0.034
Current smokers (%)	18.0	18.7	17.2	18.1	0.885
T2DM duration (yrs)*	10 (4-17)	9 (3-15)	10 (5-19)	10 (5-19)	0.007
Hba1c (%)	7.7±1.3	7.9±1.6	7.8±1.3	7.6±1.2	<0.001
BMI (kg/m <sup>2</sup> )	27.6±5.3	29.6±5.8	28.5±5.3	26.3±4.7	<0.001
Waist circumference (cm)	96.4±15.6	100.5±20.7	100.7±14.8	92.9±12.5	<0.001
SBP (mmHg)	139.4±18.3	141.0±18.3	136.0±17.7	140.3±18.3	0.001
DBP (mmHg)	78.6±8.9	79.4±9.2	79.2±9.0	78.0±8.7	0.06
IADSBP (mmHg)*	4 (2-8)	6 (3-10)	4 (2-8)	4 (2-7)	0.001
IADSBP ≥10 mmHg (%)	20.8	27.4	22.4	17.4	0.002
Heart rate (bpm)	70.9±10.9	72.2±11.6	71.0±10.3	70.3±10.8	0.049
HDL-C (mM)	1.3±0.4	1.3±0.3	1.2±0.4	1.3±0.4	<0.001
LDL-C (mM)	2.7±0.8	2.9±0.9	2.7±0.8	2.7±0.8	0.001
TG (mM)*	1.38 (1.02-1.91)	1.58 (1.11-2.07)	1.26 (1.02-1.78)	1.36 (0.99-1.89)	0.017
eGFR (ml/min/1.73 m <sup>2</sup> )*	85.6 (65.3-104.3)	82.7 (57.8-101.9)	89.3 (72.0-105.8)	84.9 (64.6-140.3)	0.007
ACR (mg/g)*	21 (6-89)	31.5 (9.9-129.3)	15.0 (4-43)	22 (5-96)	0.007
PWV (m/s)	9.7±2.9	10.1±3.1	9.5±3.2	9.6±2.8	0.122
sRAGE(pg/ml)*	693.8 (508.9-949.2)	733.6 (535.8-974.2)	651.5 (478.1-904.8)	714.1 (524.4-950.9)	0.021
Proliferative retinopathy (%)	10.7	10.1	9.3	11.6	0.201
Anti-hypertension Medications					
Usage of ACE inhibitors (%)	36.2	45.4	35.7	32.8	0.002
Usage of ARBs (%)	26.4	21.1	20.1	31.7	<0.001
CVD					
Ischemic heart disease (%)	10	8.3	14.8	8.4	0.014
History of myocardial infarction (%)	4.2	3.7	7.9	2.6	0.002
Coronary balloon angioplasty (%)	5.5	3.7	11.0	3.6	<0.001
Coronary bypass (%)	5.1	4.6	5.1	5.2	0.935
Stroke (%)	3.5	2.8	3.1	3.9	0.695

BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; HbA1c, hemoglobin A1c; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol; TG, total triglycerides; eGFR, estimated glomerular filtration rate; ACR, albumin-to-creatinine ratio ACR; PWV, pulse wave velocity; sRAGE, soluble receptor for advanced glycation end products; IADSBP, inter-arm difference in SBP; ACE, angiotensin-converting-enzyme; ARBs, angiotensin receptor blockers  
\*expressed as median (inter-quartile range)

**Table 2.** Variables associated with IADSBP  $\geq 10$  mmHg in multivariable logistic regression models in individuals with T2DM (n=1,233)

Variables	OR	(95% CI)	p value
Entry age (yrs)	1.009	(0.991-1.028)	0.318
Sex			
Male	Ref.		
Female	1.260	(0.904-1.753)	0.173
Ethnicity			
Chinese	Ref.		
Malay	1.648	(1.138-2.400)	0.009
Indian	1.380	(0.960-1.983)	0.082
Duration of diabetes (yrs)	1.006	(-0.988-1.025)	0.493
HbA1c (%)	0.985	(0.874-1.110)	0.806
BMI (kg/m <sup>2</sup> )	1.054	(1.022-1.087)	0.001
Heart rate (bpm)	1.013	(0.999-1.027)	0.077
PWV (m/s) <sup>1</sup>	0.994	(0.940-1.051)	0.834
Ln eGFR (ml/min)	1.028	(0.689-1.529)	0.893
LnACR ( $\mu\text{g}/\text{mg}$ ) <sup>2</sup>	0.978	(0.900-1.063)	0.603
Ln sRAGE per SD (pg/ml)	0.867	(0.744-1.009)	0.066
Current smoker	1.076	(0.705-1.641)	0.734
Hypertension	2.529	(1.811-3.533)	<0.001
Usage of ACE inhibitors <sup>3</sup>	1.004	(0.714-1.413)	0.980
Usage of ARBs <sup>4</sup>	1.015	(0.694-1.484)	0.939

T2DM, type 2 diabetes; HbA1c, hemoglobin A1c; BMI, body mass index; PWV, pulse wave velocity; eGFR, estimated glomerular filtration rate; ACR, albumin-to-creatinine ratio; sRAGE, soluble receptor for advanced glycation end products; ACE, angiotensin-converting-enzyme; ARBs, angiotensin receptor blockers

<sup>1</sup>Results are estimated on 1,220 subjects because of 13 (1.05%) missing values

<sup>2</sup>Results are estimated on 1,222 subjects because of 11 (0.89%) missing values

<sup>3</sup>Results are estimated on 1,228 subjects because of 5 (0.40%) missing values

<sup>4</sup>Results are estimated on 1,227 subjects because of 6 (0.49%) missing values

## DISCUSSION

In the multi-ethnic T2DM Asian cohort, we have, for the first time, demonstrated that while Malays were likely to have IADSBP  $\geq 10$  mmHg than the Chinese, the Indians have comparable odds with the Chinese. BMI and hypertension were strongly associated with IADSBP  $\geq 10$  mmHg.

The pattern of higher IADSBP in Malays and Indians compared to Chinese in this study is concordant with our previous observation of ethnic-associated greater arterial stiffness<sup>12</sup> and poor T2DM-related outcome.<sup>10,11</sup> Our finding corroborates well with prevailing literature suggesting the existence of a relationship between ethnicity and IADSBP. The IADSBP  $\geq 10$  mmHg in Chinese (6.4%,  $< 4$  mmHg),<sup>19</sup> Japanese (6.3%, 3.7 mmHg)<sup>20</sup> and Taiwanese (7.1%)<sup>21</sup> are lower than that in Caucasians in UK (8.6%, mean=4.6 mmHg,<sup>18</sup> 20%, 6.3 mmHg<sup>22</sup>) and USA (9.8%)<sup>23</sup>. In one multi-ethnic study in USA, different prevalence rates of IADSBP  $\geq 15$  mmHg were found in African-Americans (7.4%), Caucasians (5.1%), and Chinese (1.0%).<sup>24</sup> The generally lower IADSBP in Asians than Caucasians might be related to their lower BMI.<sup>20</sup> Consistently, in our study, Chinese had lower BMI, waist circumference and better lipid profile than Malays and Indians. The mechanisms underlying ethnic-disparity in vascular health is complex and may be a composite outcome of non-biological (e.g., social and economic status) and biological factors (e.g., central hemodynamics and arterial stiffness).<sup>34,35</sup>

We observed that BMI is an independent predictor of IADSBP  $\geq 10$  mmHg, which is consistent with previous

studies, suggesting a potential causal relationship between obesity and significant IADSBP.<sup>21,30,32</sup> The association between hypertension and IADSBP  $\geq 10$  mmHg is in agreement with some reports,<sup>21,30</sup> but not others.<sup>32</sup> The differences in age, SBP, DBP and BP measurement methods may all contribute to the disparate findings. Taken together, weight and BP management may ameliorate IADSBP and hence, CVD in T2DM. The odds for IADSBP  $\geq 10$  mmHg were decreased by 13.3% with 1-SD increase in Ln sRAGE. Advanced glycation end products (AGEs) play an important role in diabetic complications through interaction with the AGE receptor (RAGE). sRAGE is a multi-ligand receptor on vascular cells that acts as decoy, thus decreasing AGE-cellular binding. Experimental studies have demonstrated the role of AGE-RAGE axis in hypertension.<sup>31</sup> Decreased sRAGE level was found in non-diabetic hypertensive patients.<sup>31</sup> Therefore, further study on sRAGE and IADSBP is desirable.

The usual relative concordance of BP in both arms has been speculated to reflect on-going delicate and finely-tuned homeostatic processes. Reasons contributing to the perturbation of this homeostasis include anatomical (e.g., mechanical obstruction secondary to subclavian stenosis) and hemodynamic factors (e.g., endothelial function and atherosclerosis), which could in-part lead to an increase in IADSBP.<sup>32</sup>

CVD is known as a complex disease and can take many forms, such as stroke, myocardial infarction (MI) and ischemic heart disease (IHD).<sup>36</sup> The underlying mechanisms vary depending on the form of the disease. In our study, we found comparable occurrence of IHD, coronary balloon angioplasty and stroke in Malay and Chinese. MI in Malays is higher than Chinese, which agrees with our recent report.<sup>11</sup> It is possible IADSBP is one contributing risk factor for certain forms of CVD, such as MI. In addition to IADSBP, the development of CVD involves many different risk factors, such as genetic factors, hypertension, physical inactivity and unhealthy lifestyle. The measured and unmeasured factors may collectively explain the higher prevalence of IADSBP  $> 10$  mmHg in Malays but comparable CVD occurrence compared with Chinese.

The strength of our study is the relatively large sample size of multi-ethnic patients with T2DM living in Singapore. To the best of our knowledge, this is the first study on the association of ethnicity with IADSBP in Asians with T2DM. We are also cognizant of several limitations. We used BP measured sequentially to define IADSBP  $\geq 10$  mmHg. This might account for the prevalence of IADSBP  $\geq 10$  mmHg (20.8%) higher than most studies in T2DM.<sup>16-18</sup> The overestimation of IADSBP could also be attributable to white-coat effects and short-term large BP variability that escaped detection when using sequential measurement method.<sup>14,15</sup> Therefore, such simple sequential measurement method is recommended for screening in



routine practice, while simultaneous measurements should be reserved for confirmation.<sup>17</sup> The cross-sectional design of our study precludes any causal-inference between vascular risk factors and IADSBP. Although controlling for potential confounders, we cannot rule out residual confounding from unmeasured factors that could affect blood pressure or IADSBP across three ethnic groups, such as other classes of hypertension medications (i.e., calcium channel blockers) and other renal vascular risk factors (i.e., physical activity, carotid artery intima-media thickness and coronary artery calcium score).<sup>24,37</sup>

## CONCLUSIONS

In conclusion, our study provides the first evidence that IADSBP  $\geq 10$  mmHg in Malays with T2DM may explain their excess adverse diabetes-related vascular outcomes compared to Chinese. Therefore, measuring bilateral BP may be a simple yet clinically useful procedure to identify T2DM patients at high risk for CVD, thereby informing the need to intensify risk factor management.

## Acknowledgements

We thank Dr. Darren E. J. Seah for his assistance in subject recruitment. We also thank staff from the Singapore Clinical Research Institute (SCRI) for their contribution to study protocol and database design. Study data were collected and managed using REDCap electronic data capture tools hosted at SCRI.

## Statement of Authorship

All authors have given approval to the final version submitted.

## Author Disclosure

The authors declare that they have no competing interests.

## Funding Source

This work was supported by Singapore National Medical Research Council Grant PPG/AH(KTPH)/2011. The funder has no role in study design, data collection, analysis, interpretation and manuscript writing.

## References

- Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diab Res Clin Pract.* 2010;87(1):4-14. <https://doi.org/10.1016/j.diabres.2009.10.007>.
- Ramachandran A, Snehalatha C, Shetty AS, Nanditha A. Trends in prevalence of diabetes in Asian countries. *World J Diabetes.* 2012;3(6):110-7.
- Phan TP, Alkema L, Tai ES, Tan KHX, Yang Q, Lim WY, et al. Forecasting the burden of type 2 diabetes in Singapore using a demographic epidemiological model of Singapore. *BMJ Open Diabetes Res Care.* 2014;2(1):e000012. <https://doi.org/10.1136/bmjdr-2013-000012>.
- Jha AK, Varosy PD, Kanaya AM, et al. Differences in medical care and disease outcomes among black and white women with heart disease. *Circulation.* 2003;108(9):1089-94. <https://doi.org/10.1161/01.CIR.0000085994.38132.E5>.
- Keil JE, Sutherland SE, Knapp RG, Lackland DT, Gazes PC, Tyroler HA. Mortality rates and risk factors for coronary disease in black as compared with white men and women. *The New England Journal of Medicine.* 1993;329(2):73-8. <https://doi.org/10.1056/NEJM.199307083290201>.
- Yeo KK, Tai BC, Heng D, Lee J, Hughes K, Chew SK, et al. Ethnicity modifies the association between diabetes mellitus and ischaemic heart disease in Chinese, Malays and Asian Indians living in Singapore. *Diabetologia.* 2006;49(12):2866-73. <https://doi.org/10.1007/s00125-006-0469-z>.
- Hong CY, Chia KS, Hughes K, Ling SL. Ethnic differences among Chinese, Malay and Indian patients with type 2 diabetes mellitus in Singapore. *Singapore Med J.* 2004;45(4):154-60. PMID: 15094983.
- Dalan R, Jong M, Choo R, Chew DE, Leow MKS. Predictors of cardiovascular complication in patients with diabetes mellitus: A 5-year follow-up study in a multiethnic population of Singapore: CREDENCE II study. *Int J Cardiol.* 2013;169(4):e67-9. <https://doi.org/10.1016/j.ijcard.2013.08.128>.
- Ma S, Cutter J, Tan CE, Chew SK, Tai ES. Associations of diabetes mellitus and ethnicity with mortality in a multiethnic Asian population: Data from the 1992 Singapore National Health Survey. *Am J Epidemiol.* 2003;158(6):543-52. <https://doi.org/10.1093/aje/kwg199>.
- Liu J, Lim S, Yeoh L, et al. Long term follow-up of cardiovascular outcome in multiethnic Asians with type 2 diabetes mellitus (abstract). *Singapore Med J.* 2014;55(Suppl 1):s29.
- Low S, Chi LS, Yeoh LY, Liu JJ, Fun S, Su C, et al. Long-term diabetes outcomes in multi-ethnic Asians living in Singapore. *Diab Res Clin Pract.* 2016;111:83-92. <https://doi.org/10.1016/j.diabres.2015.09.019>.
- Zhang X, Liu JJ, Sum CF, Ying YL, Tavintharan S, Ng XW, et al. Ethnic disparity in central arterial stiffness and its determinants among Asians with type 2 diabetes. *Atherosclerosis.* 2015;242(1):22-8. <https://doi.org/10.1016/j.atherosclerosis.2015.06.019>.
- Hypertension: The Clinical Management of Primary Hypertension in Adults: Update of Clinical Guidelines 18 and 34. London, UK: Royal College of Physicians, 2011. PMID: 22855971.
- Clark CE, Taylor RS, Shore AC, Ukoumunne OC, Campbell JL. Association of a difference in systolic blood pressure between arms with vascular disease and mortality: A systematic review and meta-analysis. *Lancet.* 2012;379(9819):905-14. [https://doi.org/10.1016/S0140-6736\(11\)61710-8](https://doi.org/10.1016/S0140-6736(11)61710-8).
- Clark CE, Aboyans V. Interarm blood pressure difference: More than an epiphenomenon. *Nephrol, Dial Transplant.* 2015;30(5):695-7. <https://doi.org/10.1093/ndt/gfv075>.
- Kleefstra N, Houweling ST, Meyboom-de Jong B, Bilo HJ. [Measuring the blood pressure in both arms is of little use; longitudinal study into blood pressure differences between both arms and its reproducibility in patients with diabetes mellitus type 2]. *Ned Tijdschr Geneeskd.* 2007;151(27):1509-14. PMID: 17763810.
- Clark CE, Greaves CJ, Evans PH, Dickens A, Campbell JL. Inter-arm blood pressure difference in type 2 diabetes: A barrier to effective management? *Br J Gen Pract.* 2009;59(563):428-32. <https://doi.org/10.3399/bjgp09X420752>.
- Clark CE, Steele AM, Taylor RS, Shore AC, Ukoumunne OC, Campbell JL. Interarm blood pressure difference in people with diabetes: Measurement and vascular and mortality implications. *Diabetes Care.* 2014;37(6):1613-20. <https://doi.org/10.2337/dc13-1576>.
- Sheng CS, Liu M, Zeng WF, Huang QF, Li Y, Wang JG. Four-limb blood pressure as predictors of mortality in elderly Chinese. *Hypertension.* 2013;61(6):1155-60. <https://doi.org/10.1161/HYPERTENSIONAHA.111.00969>.
- Tokitsu T, Yamamoto E, Hirata Y, Fujisue K, Sugamura K, Maeda H, et al. Relationship between inter-arm blood pressure differences and future cardiovascular events in coronary artery disease. *J Hypertens.* 2015;33(9):1780-9. <https://doi.org/10.1097/HJH.0000000000000616>.
- Su HM, Lin TH, Hsu PC, Chu CY, Lee WH, Chen SC, et al. Association of interarm systolic blood pressure difference with atherosclerosis and left ventricular hypertrophy. *PloS One.* 2012;7(8):e41173. <https://doi.org/10.1371/journal.pone.0041173>.
- Lane D, Beevers M, Barnes N, Bourne J, John A, Malins S, et al. Interarm differences in blood pressure: When are they clinically significant? *J Hypertens.* 2002;20(6):1089-95. PMID: 12023677.
- White J, Mortensen LH, Kivimaki M, Gale CR, Batty GD. Interarm differences in systolic blood pressure and mortality among US army veterans: Aetiological associations and risk prediction in the Vietnam Experience Study. *Eur J Prev Cardiol.* 2014;21(11):1394-1400. <https://doi.org/10.1177/2047487313496193>.
- Aboyans V, Kamineni A, Allison MA, McDermott MM, Crouse JR, Ni H, et al. The epidemiology of subclavian stenosis and its association with markers of subclinal atherosclerosis: The Multi-Ethnic Study of Atherosclerosis (MESA). *Atherosclerosis.* 2010;211(1):266-70. <https://doi.org/10.1016/j.atherosclerosis.2010.01.013>.

25. Liu JJ, Tavintharan S, Yeoh LY, Sum CF, Ng XW, Pek SLT, et al. High normal albuminuria is independently associated with aortic stiffness in patients with type 2 diabetes. *Diabetic Med.* 2014;31(10):1199-1204. <https://doi.org/10.1111/dme.12461>.
26. Rognant N, Lemoine S, Laville M, Hadj-Aissa A, Dubourg L. Performance of the chronic kidney disease epidemiology collaboration equation to estimate glomerular filtration rate in diabetic patients. *Diabetes Care.* 2011;34(6):1320-2. <https://doi.org/10.2337/dc11-0203>.
27. Johansen NB, Charles M, Vistisen D, Rasmussen S, Wiinberg N, Borch-Johnsen K, et al. Effect of intensive multifactorial treatment compared with routine care on aortic stiffness and central blood pressure among individuals with screen-detected type 2 diabetes. *Diabetes Care.* 2012;35(11):2207-14. <https://doi.org/10.2337/dc12-0176>.
28. Pickering TG. What will replace the mercury sphygmomanometer? *Blood Press Monit.* 2003;8(1):23-5. <https://doi.org/10.1097/01.mbp.0000059621.89704.15>. PMID: 12604932.
29. Hosmer DW, Lemeshow S. *Applied logistic regression.* New York: John Wiley & Sons, Inc., 2000. <https://doi.org/10.1002/0471722146>.
30. Kimura A, Hashimoto J, Watabe D, Takahashi H, Ohkubo T, Kikuya M, et al. Patient characteristics and factors associated with inter-arm difference of blood pressure measurements in a general population in Ohasama, Japan. *J Hypertens.* 2004;22(12):2277-83. PMID: 15614021.
31. Geroldi D, Falcone C, Emanuele E, D'Angelo A, Calcagnino M, Buzzzi MP, et al. Decreased plasma levels of soluble receptor for advanced glycation end-products in patients with essential hypertension. *J Hypertens.* 2005;23(9):1725-9. PMID: 16093918.
32. Canepa M, Milanese Y, Ameri P, AlGhatrif M, Leoncini G, Spallarossa P, et al. Relationship between inter-arm difference in systolic blood pressure and arterial stiffness in community-dwelling older adults. *Journal of Clinical Hypertension.* 2013;15(12):880-7. <https://doi.org/10.1111/jch.12178>.
33. Report of Inter-Society Commission for Heart Disease Resources. III. Cardiovascular Disease—long-term care. Guidelines for the detection, diagnosis and management of hypertensive populations. *Circulation.* 1971;44(Suppl 5):A263-72. PMID: 5000456.
34. Thurston RC, Matthews KA. Racial and socioeconomic disparities in arterial stiffness and intima media thickness among adolescents. *Social Sci Med.* 2009;68(5):807-13. <https://doi.org/10.1016/j.socscimed.2008.12.029>.
35. Chirinos JA, Kips JG, Roman MJ, Medina-Lezama J, Li Y, Woodiwiss AJ, et al. Ethnic differences in arterial wave reflections and normative equations for augmentation index. *Hypertension.* 2011;57(6):1108-16. <https://doi.org/10.1161/HYPERTENSIONAHA.110.166348>.
36. GBD 2013 Mortality and Causes of Death Collaborators. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: A systematic analysis for the Global Burden of Disease Study 2013. *Lancet.* 2015;385(9963):117-71. [https://doi.org/10.1016/S0140-6736\(14\)61682-2](https://doi.org/10.1016/S0140-6736(14)61682-2).
37. Booth FW, Roberts CK, Laye MJ. Lack of exercise is a major cause of chronic diseases. *Compr Physiol.* 2012;2(2):1143-1211. <https://doi.org/10.1002/cphy.c110025>.

Authors are required to accomplish, sign and submit scanned copies of the JAFES Author Form consisting of: (1) the Authorship Certification that the manuscript has been read and approved by all authors, and that the requirements for authorship have been met by each author, (2) the Author Declaration that the article represents original material that is not being considered for publication or has not been published or accepted for publication elsewhere, (3) the Statement of Copyright Transfer [accepted manuscripts become the permanent property of the JAFES and are licensed with an Attribution-Share Alike-Non-Commercial Creative Commons License. Articles may be shared and adapted for non-commercial purposes as long as they are properly cited], (4) the Statement of Disclosure that there are no financial or other relationships that might lead to a conflict of interest. For Original Articles involving human participants, authors are required to submit a scanned copy of the Ethics Review Approval of their research. For manuscripts reporting data from studies involving animals, authors are required to submit a scanned copy of the Institutional Animal Care and Use Committee approval. For Case Reports or Series, and Images in Endocrinology, consent forms are required for the publication of information about patients. Articles and any other material published in the JAFES represent the work of the author(s) and should not be construed to reflect the opinions of the Editors or the Publisher.



Experience the new JAFES.  
Visit us at [www.ASEAN-endocrinejournal.org](http://www.ASEAN-endocrinejournal.org).