

PP-D-07

CHANGES IN HbA1c AND ANTHROPOMETRY IN REAL-WORLD SETTINGS IN PATIENTS WITH TYPE 2 DIABETES RECEIVING GLP-1 RECEPTOR ANALOGUES (CHARGE AUDIT)

<https://doi.org/10.15605/jafes.038.AFES.76>

Goh Kian Guan

Hospital Tengku Ampuan Afzan, Pahang, Malaysia

INTRODUCTION

GLP-1 receptor analogue is a class of glucose-lowering agents with weight loss benefits. The magnitude of benefits may not be emulated in real-world settings compared to randomised controlled trials (RCT) due to strict inclusions and frequent follow-ups. This class of medication is costly and justification is required to support its usage. This study aims to look at real-world evidence of weight loss and HbA1c changes in patients started on GLP-1 receptor analogues.

METHODOLOGY

Patients were recruited from the Diabetes Clinic in a tertiary hospital. Inclusion criteria were patients with type 2 diabetes initiated on GLP1 receptor analogues in the year 2022. Data on anthropometry, body weight and HbA1c were captured retrospectively based on patients' clinic records and segregated into 3 monthly intervals. Data were analyzed using SPSS version 22.

RESULTS

The total number of patients recruited was 14; 12 receiving subcutaneous semaglutide and two on dulaglutide. Mean body weight reduction was higher among semaglutide recipients compared to dulaglutide at 3 months, 6 months and peaked at 9 months (semaglutide -10.4% vs dulaglutide -6.0%). HbA1c changes were highest at 3 months post initiation for semaglutide (10.2% reduction from baseline), plateaued at 6 months and increased close to baseline at 9 months. There was insufficient data for dulaglutide users.

CONCLUSION

In Malaysia, the cost-per-patient-per-month for GLP1 RA is about US \$110 and about US \$10 per-patient-month for SGLT2 inhibitors, in public hospital settings. Both GLP-1 receptor analogues showed weight loss benefits, with the effect seen greater in subcutaneous semaglutide users. This finding has been reproduced in other studies that have shown more weight loss effects when switching from dulaglutide to semaglutide. Moreover, the weight loss effect is continuous up to the 9th month for both drugs and did not plateau. Despite the further weight loss, the HbA1c effect was maximum at 3rd month and plateaued

at 6 months onwards. Our study findings were comparable with other existing real-world evidence and RCTs, in line with the latest Malaysian guidelines in managing patients with type 2 diabetes and excessive weight. The limitations of this study were the small sample size and more subjects received semaglutide.

KEYWORDS

semaglutide, dulaglutide, obesity, type 2 diabetes, GLP-1 receptor analogues

PP-D-08

THE EFFICACY OF TELEMONITORING AND INTEGRATED PERSONALIZED DIABETES MANAGEMENT (IPDM) IN PEOPLE WITH INSULIN-TREATED TYPE 2 DIABETES MELLITUS: A PRELIMINARY ANALYSIS

<https://doi.org/10.15605/jafes.038.AFES.77>

Primploy Greeviroj,¹ Thanyalak Saetang,¹ Subhanudh Thavaraputtam,¹ Prangareeya Santisitthanon,¹ Natnicha Hounggam²

¹*Division of Endocrinology and Metabolism, Department of Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand)*

²*Excellence Centre for Diabetes, Hormones, and Metabolism, King Chulalongkorn Memorial Hospital, Bangkok, Thailand*

INTRODUCTION

Integrated personalized diabetes management (iPDM) can improve glycemic control in people with diabetes. Emerging evidence suggests that telehealth can improve diabetes care. The purpose of this study was to assess the efficacy of diabetes care through a structured telehealth model of care.

METHODOLOGY

A 6-month single-center, open-labeled, prospective randomized controlled trial enrolled insulin-treated subjects with diabetes, aged 18-65 years old and A1c of 7.4-10.5%. All participants received standard diabetes education. The tele-iPDM group will connect their glucometer to the cloud-based telemonitoring platform and adjust insulin following a protocol by investigators weekly for 3 months (phase 1), then monthly for another 3 months (phase 2). The usual care group will receive standard diabetes care and record glucose data in the paper logbook. The primary outcome was a difference in A1c change from baseline between 2 groups at 12 and 24 weeks. Secondary outcomes included changes in FPG, BW, BMI, and the percentage of people with A1c 0.5% at 24 weeks, SMBG profiles and the number of hypoglycemic events.

RESULTS

Sixty-one subjects completed the study. The mean age of participants was 53.07±7.74 years. The mean duration of diabetes was 11.76 ± 8.26 years. The baseline A1c was 8.48 ± 0.76%. Phase 1 study showed a mean reduction in A1c of 1.02% (95% CI: 0.74–1.30) in the tele-iPDM group and 0.48% (95% CI: 0.19–0.76) in the usual care group. The difference in A1c reduction between the 2 groups was 0.55% [95% CI: 0.15–0.95, $p < 0.05$]. At 24 weeks of follow-up, the mean difference in A1c between the tele-iPDM and usual care groups is 0.72% [95% CI: 0.24–1.20, $p < 0.05$]. There were no significant differences in body weight and body mass index and hypoglycemic events between both groups.

CONCLUSION

Telemonitoring can facilitate the iPDM care model in people with insulin-treated type 2 diabetes mellitus. It improves the efficiency of diabetes care and improves glycemic control at 12 weeks and can maintain glycemic control at 24 weeks.

KEYWORDS

telemonitoring, structured feedback loop, type 2 diabetes, insulin-treated

PP-D-09

IMPACT OF DIABETES AND SARCOPIENIA ON MORTALITY

<https://doi.org/10.15605/jafes.038.AFES.78>

Eyun Song and Kyung Mook Choi

Division of Endocrinology and Metabolism, Korea University

INTRODUCTION

Diabetes mellitus (DM) and sarcopenia are growing public health threats in an aging society. They share common pathophysiological mechanisms and are associated with serious health consequences. We investigated the impact of DM and sarcopenia on all-cause and cardiovascular mortalities in a longitudinal nationwide population-based study.

METHODOLOGY

Subjects who participated in the Korea National Health and Nutrition Examination Survey conducted from 2008 to 2011 with available appendicular skeletal muscle mass data were analyzed. Mortality data up to December 2020 were retrieved from the National Death Registry.

RESULTS

Of the 17,920 subjects, 14,737 (82.2%) had neither DM nor sarcopenia (DM-/SP-), 1,349 (7.5%) had only DM (DM+/SP-), 1,425 (8.0%) had only sarcopenia (DM-/SP+), and 409 (2.3%) had both DM and sarcopenia (DM+/SP+). Compared to the DM-/SP- group, all-cause mortality was increased,

with hazard ratios (HRs) of 1.29 (95% confidence interval [CI]: 0.97–1.73), 1.44 (95% CI: 1.12–1.85), and 1.88 (95% CI: 1.29–2.73) in the DM+/SP-, DM-/SP+, and DM+/SP+ groups, respectively, after adjusting for covariates. The data showed the DM+/SP+ group had the highest risk of overall mortality (p -for-trend = 0.042). Cardiovascular mortality was increased, with HRs of 1.34 (95% CI: 0.79–2.25), 1.39 (95% CI: 0.82–2.36), and 1.98 (95% CI: 1.04–3.77) in the DM+/SP-, DM-/SP+ and DM+/SP+ groups, respectively, compared to DM-/SP- group (p -for-trend 0.037).

CONCLUSION

The coexistence of DM and sarcopenia synergistically increased the risk of all-cause and cardiovascular mortality. Individuals with either disease may require more careful management to prevent the development of the other disease to reduce mortality.

KEYWORDS

diabetes, sarcopenia, mortality

PP-D-10

EFFECTIVENESS OF A DIABETES ONE-STOP CLINIC FOR TYPE 2 DIABETES PATIENTS IN A TERTIARY CARE HOSPITAL IN THAILAND

<https://doi.org/10.15605/jafes.038.AFES.79>

Jintanan Jangsiripornpakorn,¹ Phawinpon Chotwanvirat,¹ Nuttha Khunthupat,² Sumitra Chukaew¹

¹*Diabetes and Metabolic Care Center, Taksin Hospital, Bangkok, Thailand*

²*Taksin Hospital, Bangkok, Thailand*

INTRODUCTION

A multidisciplinary team approach is a strategy for optimizing care for patients with uncontrolled type 2 diabetes. However, the effectiveness of an integrated diabetes care team remains unclear.

METHODOLOGY

This study aims to evaluate the efficacy of a diabetes one-stop clinic, a multidisciplinary outpatient clinic that aims to provide integrated care for type 2 diabetes at Taksin Hospital. A multidisciplinary team, consisting of an endocrinologist, a certified nurse educator, a pharmacist, and a nutritionist, attended a weekly clinic at the Diabetes and Metabolic Care Center in Taksin Hospital. The integrated care team provides diabetes self-management and support, nutritional counseling, and diabetes management. To evaluate the change in their metabolic profile, a retrospective assessment of medical records for type 2 diabetes patients who visited a clinic between October 2021 and March 2022 and had HbA1c above 8%