



CONCLUSION

These results suggest that the genotyping of cf-DNA of APA from AVS samples is promising to detect the somatic mutations present in the APA. However, as AVS is an invasive procedure, genotyping of cf-DNA from peripheral blood may be investigated as an alternative. Therefore, further work is needed to ensure this strategy can be non-invasive as then it can be used as a screening method before AVS.

PP-A-05

ELUCIDATING THE EFFECTS OF MUTATIONS IN Q209 OF GNA11 ON CELL APOPTOSIS IN HUMAN ADRENOCORTICAL CELLS

<https://doi.org/10.15605/jafes.037.AFES.23>

Kha Chin Long,¹ **Norlela Sukor**,¹ **Geok Chin Tan**,² **Nor Azian Abdul Murad**,³ **Morris Jonathan Brown**,⁴ **Elena Aisha Azizan**¹

¹The National University of Malaysia (UKM) Medical Centre, Cheras, Malaysia

²The National University of Malaysia (UKM), Cheras, Malaysia

³UKM Medical Molecular Biology Institute, Cheras, Malaysia

⁴Queen Mary University of London, London, United Kingdom

OBJECTIVES

Gain-of-function mutations, Q209H, Q209P and Q209L, of GNA11 were recently found to occur in CTNNB1 mutant aldosterone-producing adenomas (APAs). These mutations were also found to be present in the hyperplastic zona glomerulosa adjacent to the double-mutant APAs. This study aims to investigate the effects of GNA11 Q209 mutations on tumorigenesis through measurement of cell apoptosis in the human adrenocortical cell line, HAC15.

METHODOLOGY

HAC15 was transfected with GFP-tagged GNA11 Q209H, Q209L, Q209P or wild-type (WT) plasmids. To note, HAC15, a subclone of H295R cells, inherently has the CTNNB1 S45P mutation. 48 hours post-transfection, cell apoptosis was assessed using the Pacific Blue™ Annexin V/SYTOX™ AADvanced™ apoptosis assay (BD Biosciences, USA). The supernatants and cells were harvested for aldosterone and cortisol determination, and RNA isolation.

RESULTS

HAC15 cells transfected with GNA11 mutants, Q209H, Q209L and Q209P, had elevated aldosterone production compared to WT at 62.4% (p=0.001), 71.2% (p=0.001) and 59.5% (p=0.001), respectively. Cortisol production was only slightly elevated in HAC15 cells transfected with Q209H (19.7%, p=0.01) and Q209L (24.6%, p=0.01), compared to WT. CYP11B2 mRNA expression was also upregulated compared to WT by 3.5 folds (p=0.001) for Q209H, and around 8 folds (p=0.001) for Q209L and Q209P. Analysis of flow cytometric apoptosis assay showed GNA11 mutants did not affect cell apoptosis.

CONCLUSION

The findings suggests that GNA11 Q209 mutation increases aldosterone secretion of adrenocortical cells with no or little effect on apoptosis rate. Further experiments on cell proliferation are needed to rule out whether GNA11 Q209 mutations affects tumorigenesis.

PP-A-06

UTILITY OF ADRENAL VENOUS SAMPLING IN ACTH-INDEPENDENT CUSHING'S SYNDROME PRESENTING WITH BILATERAL ADRENAL ADENOMA

<https://doi.org/10.15605/jafes.037.AFES.24>

Pamela Ann Aribon and **Rowena Ejercito-De Jesus**
St. Luke's Medical Center Global City, Taguig, Philippines

BACKGROUND

Adrenocorticotrophic Hormone (ACTH)-independent Cushing's syndrome in a patient with bilateral adenoma poses a management challenge to clinicians. Utilization of adrenal venous sampling (AVS), as in this case, is instrumental in the precise localization of the functioning adenoma which will ensure the best management for these patients.

CASE

We report the case of a 67-year-old Filipino who presented with gradual weight gain for 3 months described as rounding of the face and increasing abdominal girth. The diagnosis of ACTH-independent Cushing's syndrome was based on undetectable ACTH and an unsuppressed cortisol level by dexamethasone suppression test. CT scan revealed bilateral adrenal adenomas measuring 1.1 x 0.9 cm (APxT) in the right and 1.1 x 1.3 cm (APxT) in the left. AVS was done using cortisol levels adjusted by plasma aldosterone. This successfully lateralized the hypersecretion of cortisol to the left adrenal gland, hence a unilateral laparoscopic left adrenalectomy was done.