

Letter to the Editor on the article entitled, 'Perioperative Complications Associated with Routine Preoperative Glucocorticoid Use Among Patients Undergoing Pituitary Surgery with Normal Preoperative HPA Axis: A Retrospective Cohort Study,' by Magnaye and Paz-Pacheco, published in JAFES Vol. 40. No. 1.

Eisha Fatimah,<sup>1</sup> Sarosh Usman,<sup>1</sup> Muhammad Aqib Farooq Khan<sup>2</sup>

<sup>1</sup>Quaid-e-Azam Medical College, Bahawalpur, Pakistan

<sup>2</sup>Islamic International Medical College, Riphah International University, Rawalpindi, Pakistan

**Key words:** hypothalamic-pituitary-adrenal axis, glucocorticoids, perioperative complications

This letter addresses the recent study by Magnaye et al., which investigated the impact of routine preoperative glucocorticoid administration in patients with a normal hypothalamic-pituitary-adrenal (HPA) axis undergoing pituitary surgery. The authors are commended for their valuable contribution to understanding steroid-related perioperative outcomes. However, we highlight the need for further clarification regarding the wide range of steroid regimens reported and the absence of detailed information on timing, indications, and factors influencing steroid administration. Additionally, differentiating between transient and permanent postoperative diabetes insipidus could provide deeper insight into the observed higher incidence among steroid-treated patients. Addressing these aspects, along with establishing standardized steroid protocols, could enhance the specificity and clinical applicability of the study's findings.

I recently came across this study by Magnaye et al.,<sup>1</sup> in which the authors have done a commendable job in addressing a clinically important topic, i.e., whether the routine administration of preoperative glucocorticoids in patients undergoing pituitary surgery with a normal hypothalamic-pituitary-adrenal (HPA) axis is actually beneficial in decreasing the rate of perioperative complications such as diabetes insipidus and postoperative infections. The study has done a great job of comparing the effects of steroid vs. non-steroid administration. The important findings highlighted by the authors provide significant and valuable points that shed light on this topic in a very detailed way.

The authors' work is commendable. However, we would like to bring to the authors' attention to some points. The authors have provided a wide range of regimens (mean hydrocortisone equivalent  $\approx$  142 mg/day, range 50–583 mg/

day). In this case, a proper breakdown about timing and any additional indications or factors influencing the decision of steroid administration could have better explained patients having a higher risk of developing complications.<sup>2</sup> Also, the authors have reported a higher incidence of postoperative diabetes insipidus in patients given preoperative steroids (52.5% vs 28.2%,  $p = 0.006$ ), but specifying the type of diabetes insipidus (permanent or transient) could maybe clear some doubts regarding the underlying cause. Transient diabetes insipidus is often seen after pituitary surgery and could have other causative factors, so the differentiation could have led to other potential causes being identified.<sup>3</sup>

In summary, this study by Magnaye et al., has raised valid concerns regarding the increased risk of certain postoperative infections in patients given steroids preoperatively. It is truly commendable and provides a good insight into avoiding complications that could arise as a result of unnecessary steroid administration. However, mentioning the above points could add a touch of more specificity to the study. Discussing standardized steroid protocols that support the steroid sparing strategy and specifying the type of diabetes insipidus would help enhance the understanding of outcomes.

#### References

1. Magnaye FM, Paz-Pacheco E. Perioperative complications associated with routine preoperative glucocorticoid use among patients undergoing pituitary surgery with normal preoperative HPA axis: A retrospective cohort study. *ASEAN Fed Endocr Soc.* 2025;40(1):89-96. PMID: 40416483 PMCID: PMC12097979 DOI: 10.15605/jafes.040.01.06
2. Chen Cardenas SM, Santhanam P, Morris-Wiseman L, et al. Perioperative evaluation and management of patients on glucocorticoids. *J Endocr Soc.* 2022;7(2):bvac185. PMID: 36545644 PMCID: PMC9760550 DOI: 10.1210/jendso/bvac185
3. Almalki MH, Ahmad MM, Brema I, et al. Management of diabetes insipidus following surgery for pituitary and suprasellar tumours. *Sultan Qaboos Univ Med J.* 2021;21(3):354-64. PMID: 34522399 PMCID: PMC8407907 DOI: 10.18295/squmj.4.2021.010

## RESPONSE OF THE AUTHORS

The authors appreciate the questions raised in relation to the paper “Perioperative complications associated with routine preoperative glucocorticoid use among patients undergoing pituitary surgery with normal preoperative HPA axis: a retrospective cohort study.”

We are pleased to know that readers value the issues related to this important clinical condition. In response to the points raised, indeed, these are critical factors that can provide more clarity to the relationship between preoperative steroid use and perioperative outcomes. As described, this was a retrospective study that relied on available data on patients’ medical records.

Ascertainment of detailed information on timing, indications and other factors determining choice for steroid administration was limited. Determination of transient versus permanent diabetes insipidus was not specified. Thus, as stated in the paper, it is recommended that a prospective study be carried out with a randomized controlled trial (RCT) design in a larger group of patients. Thereafter, a consensus can be developed and further discussed.

**Franz Michael Magnaye and Elizabeth Paz-Pacheco**

Authors are required to accomplish, sign and submit scanned copies of the JAFES Author Form consisting of: (1) Authorship Certification, that authors contributed substantially to the work, that the manuscript has been read and approved by all authors, and that the requirements for authorship have been met by each author; (2) the Author Declaration, that the article represents original material that is not being considered for publication or has not been published or accepted for publication elsewhere, that the article does not infringe or violate any copyrights or intellectual property rights; that no references have been made to predatory/suspected predatory journals; and that use of artificial intelligence (AI) or AI-assisted technologies shall be declared to include the name of the AI tool or service used; (3) the Author Contribution Disclosure, which lists the specific contributions of authors; (4) the Author Publishing Agreement which retains author copyright, grants publishing and distribution rights to JAFES, and allows JAFES to apply and enforce an Attribution-Non-Commercial Creative Commons user license; and (5) the Conversion to Visual Abstracts (\* optional for original articles only) to improve dissemination to practitioners and lay readers. Authors are also required to accomplish, sign, and submit the signed ICMJE form for Disclosure of Potential Conflicts of Interest. For original articles, authors are required to submit a scanned copy of the Ethics Review Approval of their research as well as registration in trial registries as appropriate. For manuscripts reporting data from studies involving animals, authors are required to submit a scanned copy of the Institutional Animal Care and Use Committee approval. For Case Reports or Series, and Images in Endocrinology, consent forms, are required for the publication of information about patients; otherwise, appropriate ethical clearance has been obtained from the institutional review board. Articles and any other material published in the JAFES represent the work of the author(s) and should not be construed to reflect the opinions of the Editors or the Publisher.



**Send your paper to the publication pathway.  
Instructions to Authors at  
[www.ASEAN-endocrinejournal.org](http://www.ASEAN-endocrinejournal.org).**